


# **Pascagoula River Phase One Total Maximum Daily Load for Mercury**



## **Pascagoula River Basin**



**Prepared By  
Mississippi Department of Environmental Quality  
TMDL/WLA Branch  
P.O. Box 10385  
Jackson, MS 39289-0385  
(601) 961-5171  
[www.deq.state.ms.us](http://www.deq.state.ms.us)**



## FOREWORD

This report has been prepared in accordance with the schedule contained within the federal consent decree dated December 22, 1998 (*Sierra Club v. Hankinson, No. 97-CV-3683 (N.D. Ga.)*). The report contains one or more Total Maximum Daily Loads (TMDLs) for water body segments found on Mississippi's 1996 Section 303(d) List of Impaired Waterbodies. Because of the accelerated schedule required by the consent decree, many of these TMDLs have been prepared out of sequence with the State's rotating basin approach. The segments addressed are comprised of monitored segments that have data indicating impairment. The implementation of the TMDLs contained herein will be prioritized within Mississippi's rotating basin approach.

The amount and quality of the data on which this report is based are limited. As additional information becomes available, the TMDLs may be updated. Such additional information may include water quality and quantity data, changes in pollutant loadings, modification to state water quality criteria, or changes in landuse within the watershed. In some cases, additional water quality data may indicate that no impairment exists.

### Prefixes for fractions and multiples of SI units

| Fraction   | Prefix | Symbol | Multiple  | Prefix | Symbol |
|------------|--------|--------|-----------|--------|--------|
| $10^{-1}$  | deci   | d      | 10        | deka   | da     |
| $10^{-2}$  | centi  | c      | $10^2$    | hecto  | h      |
| $10^{-3}$  | milli  | m      | $10^3$    | kilo   | k      |
| $10^{-6}$  | micro  | $\mu$  | $10^6$    | mega   | M      |
| $10^{-9}$  | nano   | n      | $10^9$    | giga   | G      |
| $10^{-12}$ | pico   | p      | $10^{12}$ | tera   | T      |
| $10^{-15}$ | femto  | f      | $10^{15}$ | peta   | P      |
| $10^{-18}$ | atto   | a      | $10^{18}$ | exa    | E      |

### Conversion Factors

| To convert from | To        | Multiply by | To Convert from              | To       | Multiply by |
|-----------------|-----------|-------------|------------------------------|----------|-------------|
| acres           | sq. miles | 0.001562    | days                         | seconds  | 86400.00    |
| cubic feet      | cu. Meter | 0.028316    | feet                         | meters   | 0.304800    |
| cubic feet      | gallons   | 7.480519    | gallons                      | cu. feet | 0.133680    |
| cubic feet      | liters    | 28.31684    | hectares                     | acres    | 2.471053    |
| cfs             | gal/min   | 448.8311    | miles                        | meters   | 1609.344    |
| cfs             | MGD       | 0.646316    | mg/l                         | ppm      | 1           |
| cubic meters    | gallons   | 264.1720    | $\mu\text{g/l} * \text{cfs}$ | gm/day   | 2.45        |
| cubic meters    | liters    | 1000        | $\mu\text{g/l} * \text{MGD}$ | gm/day   | 3.79        |

# CONTENTS

|  | <u>Page</u> |
|--|-------------|
| TMDL INFORMATION PAGE.....                                       | vi          |
| EXECUTIVE SUMMARY .....  | viii        |
| 1.0 INTRODUCTION .....   | 1           |
| 1.1 BACKGROUND .....   | 1           |
| 1.2 PHASED TMDL APPROACH .....                                   | 1           |
| 1.3 WATER BODY SEGMENT LOCATION.....                             | 1           |
| 1.4 WATER BODY DESIGNATED USE.....                               | 5           |
| 1.5 APPLICABLE WATER QUALITY STANDARDS .....                     | 5           |
| 2.0 TMDL ENDPOINT AND WATER QUALITY ASSESSMENT.....              | 6           |
| 2.1 SELECTION OF A TMDL ENDPOINT AND CRITICAL CONDITION.....     | 6           |
| 2.1.1 Mercury Speciation and Chemistry .....                     | 6           |
| 2.1.2 Mercury Transport and Transformations .....                | 7           |
| 2.1.3 Mississippi Mercury Criteria (Fresh Water).....            | 8           |
| 2.2 MISSISSIPPI REGULATIONS ON FLOW DETERMINATION .....          | 9           |
| 2.3 DISCUSSION OF INSTREAM WATER QUALITY .....                   | 9           |
| 2.3.1 Inventory of Water Quality Monitoring Data .....           | 9           |
| 2.3.2 Analysis of Fish Tissue Data .....                         | 10          |
| 3.0 SOURCE ASSESSMENT .....                                      | 11          |
| 3.1 POTENTIAL SOURCES OF MERCURY.....                            | 11          |
| 3.1.1 Fluorescent and High-Intensity Discharge Lamps.....        | 12          |
| 3.1.2 Mercury Switches and Relays.....                           | 12          |
| 3.1.3 Mercury-Containing Thermostats and Thermostat Probes ..... | 12          |
| 3.1.4 Mercury Thermometers .....                                 | 13          |
| 3.1.5 Gauges, Manometers, Barometers, and Vacuum Gauges.....     | 13          |
| 3.2 POINT SOURCE ASSESSMENT .....                                | 13          |
| 4.0 MODELING PROCEDURE .....                                     | 16          |
| 4.1 MODELING CALCULATIONS .....                                  | 16          |
| 4.2 CALCULATION SETUP.....                                       | 16          |
| 4.3 SOURCE REPRESENTATION.....                                   | 16          |

5.0 ALLOCATION..... 18

    5.1 TMDL CALCULATION..... 18

    5.2 TMDL ALLOCATIONS ..... 19

        5.2.1 Wasteload Allocations ..... 19

        5.2.2 Load Allocations..... 20

    5.3 INCORPORATION OF A MARGIN OF SAFETY ..... 21

    5.4 SEASONALITY ..... 22

    5.5 IMPLEMENTATION PLAN ..... 22

6.0 CONCLUSION..... 24

    6.1 FOLLOW-UP MONITORING..... 24

    6.2 PUBLIC PARTICIPATION ..... 24

DEFINITIONS..... 25

ABBREVIATIONS ..... 29

REFERENCES ..... 30

APPENDIX A: Fish Tissue Data..... 32

APPENDIX B: Daily Load from NPDES Permitted Facilities ..... 35

APPENDIX C: Mercury Use Outline ..... 36

APPENDIX D: Mississippi Fish Advisories ..... 40

**TABLES**

Table i. Listing Information ..... vi

Table ii. Water Quality Standard..... vi

Table iii. NPDES Facilities ..... vii

Table iv. Total Maximum Daily Load..... vii

Table 1. Water Body Identification for the Pascagoula River Phase One Mercury TMDL .....2

Table 2. Pascagoula River Watershed Landuse Distribution.....4

Table 3. State of Mississippi Water Quality Criteria for Intrastate, Interstate, and Coastal Waters .....5

Table 4. Water Quality Station Data Analysis .....10

Table 5. Permitted Facilities in the Pascagoula River Watershed.....13

Table 6. Facilities in the Pascagoula River Watershed Recommended for Mercury Monitoring.....17

Table 7. TMDL for Total Mercury II.....18

Table 8. Facilities Recommended to Perform Mercury Effluent Monitoring.....20

Table 9. Pollution Prevention (P2) Alternatives for Products Containing Mercury .....23

**FIGURES**

Figure 1. Location of the Pascagoula River Watershed .....2

Figure 2. The Pascagoula River Watershed – 303(d) Segment Locations .....3

Figure 3. Pascagoula River Watershed Landuse Distribution.....4

**TMDL INFORMATION PAGE**

**Table i. Listing Information**

| Name  | ID        | County         | HUC      | Cause   | Mon/Eval  |
|---|-----------|----------------|----------|---------|-----------|
| Pascagoula River  | MSPASRM1  | George Jackson | 03170006 | Mercury | Monitored |
| Near Benndale: From confluence with Leaf River and Chickasawhay River to Pascagoula River Split |           |                |          |         |           |
| East Pascagoula River   | MSEPASRM1 | Jackson        | 03170006 | Mercury | Monitored |
| Near Colltown: From East West Split near Barnes Lake to confluence with Escatawpa River         |           |                |          |         |           |
| East Pascagoula River   | MSEPASRM2 | Jackson        | 03170006 | Mercury | Monitored |
| Near Pascagoula: From confluence with Escatawpa River to mouth at Mississippi Sound             |           |                |          |         |           |
| West Pascagoula River   | MSWPASRM1 | Jackson        | 03170006 | Mercury | Monitored |
| Near Colltown: From East West Split to confluence with Mongers Creek                            |           |                |          |         |           |
| West Pascagoula River   | MSWPASRM2 | Jackson        | 03170006 | Mercury | Monitored |
| Near Vancleave: From confluence with Mongers Creek to mouth at Mississippi Sound                |           |                |          |         |           |

**Table ii. Water Quality Standard**

| Parameter   | Beneficial use          | Water Quality Criteria   |
|---|-------------------------|--|
| Mercury (II)<br>total dissolved Hg(II)<br>expressed as total<br>recoverable | Aquatic Life<br>Support | <p><i>Fresh Water</i></p> <p><u>Acute</u>: instantaneous concentration may not exceed 2.1 µg/l<br/> <u>Chronic</u>: average concentration may not exceed 0.012 µg/l<br/>                     expressed as total recoverable</p> <p><i>Salt Water</i></p> <p><u>Acute</u>: instantaneous concentration may not exceed 1.8 µg/l<br/> <u>Chronic</u>: average concentration may not exceed 0.025 µg/l<br/>                     expressed as total recoverable</p> |

**Table iii. NPDES Facilities**

| <b>NPDES ID</b> | <b>Facility Name</b>   | <b>County</b> | <b>Receiving Water</b>                             | <b>Flow (MGD)</b> |
|-----------------|--|---------------|--|-------------------|
| MS0031828       | George County Schools<br>Central Elementary School               | George        | Unnamed Tributary to<br>Big Creek                  | 0.0160            |
| MS0044504       | Lucedale POTW  | George        | Big Cedar Creek                                    | 0.5000            |
| MS0020249       | MS Gulf Coast Regional Wastewater<br>Authority - Pascagoula POTW | Jackson       | Pascagoula River                                   | 10.0000           |
| MS0028355       | Jackson County School District<br>East Central School            | Jackson       | Black Creek  | 0.0500            |
| MS0028762       | Jackson County School District<br>Vancleave Junior High School   | Jackson       | Unnamed Tributary to<br>Bluff Creek                | 0.0240            |
| MS0032115       | Bluff Creek Mobile Home Park                                     | Jackson       | Bluff Creek  | 0.0340            |
| MS0033375       | Kwik Kar Wash  | Jackson       | Unnamed Tributary to<br>Mary Walker Bayou          | 0.0009            |
| MS0035637       | Gulf City Seafoods, Inc.   | Jackson       | Pascagoula River                                   | 0.0297            |
| MS0035769       | MDOT Interstate 10 Rest Area                                     | Jackson       | Marsh to Farrigut Lake to<br>West Pascagoula River | 0.0500            |
| MS0038326       | Jackson County School District<br>Vancleave High School          | Jackson       | Unnamed Tributary to Bluff<br>Creek                | 0.0165            |
| MS0042269       | Finicky Pet Food, Inc.   | Jackson       | East Pascagoula River                              | 0.0040            |
| MS0043010       | MS Gulf Coast Regional Wastewater<br>Authority - Gautier POTW    | Jackson       | West Pascagoula River                              | 4.0000            |
| MS0043966       | Seachick, Inc.   | Jackson       | Unnamed Tributary to Clarke<br>Bayou               | 4.4300            |
| MS0046221       | Davis Laundry and Quickstop                                      | Jackson       | Black Creek  | 0.0036            |
| MS0047007       | Circle V Traditions, Inc.  | Jackson       | Unnamed Tributary to Little<br>Bluff Creek         | 0.0024            |
| MS0048445       | Colle Towing Company, Inc.                                       | Jackson       | Pascagoula River                                   | 0.0050            |
| MS0055379       | Pascagoula Water Treatment Plant -<br>Communy Avenue             | Jackson       | Unnamed Creek to Yazoo Lake                        | 60.4500           |

**Table iv. Total Maximum Daily Load**

| <b>WLA</b>  | <b>LA</b>    | <b>MOS</b>   | <b>Type</b>           | <b>TMDL</b> |
|-------------|--------------|--------------|-----------------------|-------------|
| 3.62 gm/day | 11.53 gm/day | 15.15 gm/day | implicit and explicit | 30.3 gm/day |

## **EXECUTIVE SUMMARY**

According to Mississippi's 2002 List of Water Bodies, portions of the Pascagoula River are impaired by mercury. The listed segments are located within Hydrologic Unit Code (HUC) 03170006 in George and Jackson Counties in south Mississippi. Largemouth bass, flathead catfish, channel catfish, bass sp., spotted bass, blue catfish, bowfin, red drum, sheepshead, white trout, Atlantic croaker, and gafftopsail caught in this water body have been sampled and the data indicate a possible impairment due to levels of mercury in the fish flesh. Based on these data, the State of Mississippi issued fish consumption advisories (see Appendix D) for portions of the Pascagoula River. These advisories were issued to help protect people who regularly consume fish caught in the water bodies. The bioaccumulation of methylmercury in fish flesh is the basis for the impairment in the water body.

This Phase One Mercury TMDL for the Pascagoula River has been developed prior to a complete understanding of the linkage between mercury in the water and mercury in the fish. Additionally, this Phase One Mercury TMDL is only concerned with point source contributions to the water body. Atmospheric deposition, nonpoint source contributions, and natural background will be considered in Phase Two. It is anticipated that the mercury data generated from the point source contributors during the next few years will enhance the knowledge base on this issue.

The endpoints selected for this Phase One Mercury TMDL are based on MDEQ regulations. There are several mercury criteria to evaluate. The human health criterion is currently 153 ng/l of total mercury. The aquatic life support criteria are 12 ng/l fresh water and 25 ng/l salt water of total mercury II expressed as total recoverable. Recent EPA criteria guidance has suggested that each of these numbers need to be revised. This guidance recommends that the 153 ng/l total mercury criterion be replaced with a methylmercury criterion of 0.3 mg/kg measured in fish tissue. The guidance also recommends that the aquatic life support numbers increase to a more representative value of 770 ng/l and 940 ng/l total mercury, respectively. However, these new numbers have not been adopted by the Mississippi Commission on Environmental Quality. MDEQ is therefore proposing the most protective of the currently adopted criteria, 12 ng/l. To further account for the unknowns, an additional explicit margin of safety is included in this TMDL. This explicit margin of safety is set at 50%.

The implementation plan in this Phase One TMDL calls for a moratorium on any future increase in mercury discharges in the Pascagoula River watershed. Increased monitoring in the Pascagoula River watershed is recommended. This TMDL also recommends pollution prevention alternatives and activities.



## **1.0 INTRODUCTION**

### **1.1 BACKGROUND**

The identification of water bodies not meeting their designated use and the development of total maximum daily loads (TMDLs) for those water bodies are required by Section 303(d) of the Clean Water Act. This is also a requirement of the Environmental Protection Agency's (EPA) Water Quality Planning and Management Regulations (40 CFR Part 130). The TMDL process is designed to restore and maintain the quality of those impaired water bodies through the establishment of pollutant specific allowable loads. The pollutant of concern for this TMDL is mercury. The purpose of this TMDL is to establish water quality objectives and best management practices to reduce the mercury levels currently found in fish flesh taken from the Pascagoula River.

Human exposure to inorganic mercury in large amounts can cause a variety of health effects. The two organ systems most likely affected are the central nervous system and the kidney. However, the most significant concerns regarding chronic exposure to low concentrations of methylmercury in fish are for neurological effects on the developing fetus and children.<sup>1</sup>

### **1.2 PHASED TMDL APPROACH**

This document is phase one of a multi-phase TMDL being developed for mercury in the Pascagoula River. This Phase One Mercury TMDL will determine the maximum load of mercury that should be introduced into the impaired segments based on Mississippi's water quality criteria. Phase Two of this TMDL project, to be completed at a later date, will quantify the mercury load to this water body that is directly related to atmospheric sources and other nonpoint sources. Phase Two will also attempt to include a fate and transport model for the water body that will better characterize aquatic mercury cycling.

### **1.3 WATER BODY SEGMENT LOCATION**

As summarized in Table 1, five segments of the Pascagoula River have been listed as impaired due to mercury in fish tissue. The location of the Pascagoula River watershed is provided in Figure 1. The location of the 303(d) listed segments are shown in Figure 2. Figure 3 and Table 2 provide landuse information for the watershed.

Table 1. Water Body Identification for the Pascagoula River Phase One Mercury TMDL

| Name  | ID        | County            | HUC      | Cause   | Mon/Eval  |
|---|-----------|-------------------|----------|---------|-----------|
| Pascagoula River  | MSPASRM1  | George<br>Jackson | 03170006 | Mercury | Monitored |
| Near Benndale: From confluence with Leaf River and Chickasawhay River to Pascagoula River Split |           |                   |          |         |           |
| East Pascagoula River   | MSEPASRM1 | Jackson           | 03170006 | Mercury | Monitored |
| Near Colltown: From East West Split near Barnes Lake to confluence with Escatawpa River         |           |                   |          |         |           |
| East Pascagoula River   | MSEPASRM2 | Jackson           | 03170006 | Mercury | Monitored |
| Near Pascagoula: From confluence with Escatawpa River to mouth at Mississippi Sound             |           |                   |          |         |           |
| West Pascagoula River   | MSWPASRM1 | Jackson           | 03170006 | Mercury | Monitored |
| Near Colltown: From East West Split to confluence with Mongers Creek                            |           |                   |          |         |           |
| West Pascagoula River   | MSWPASRM2 | Jackson           | 03170006 | Mercury | Monitored |
| Near Vanclave: From confluence with Mongers Creek to mouth at Mississippi Sound                 |           |                   |          |         |           |

Figure 1. Location of the Pascagoula River Watershed




Figure 2. The Pascagoula River Watershed – 303(d) Segment Locations




Figure 3. Pascagoula River Watershed Landuse Distribution




Table 2. Pascagoula River Watershed Landuse Distribution

|            | Urban  | Forest  | Cropland | Pasture/Grass | Disturbed | Water  | Wetlan | Total   |
|------------|--------|---------|----------|---------------|-----------|--------|--------|---------|
| Acres      | 10,452 | 164,319 | 3,085    | 48,833        | 60,579    | 13,441 | 85,386 | 386,095 |
| Percentage | 2.7%   | 42.6%   | 0.8%     | 12.6%         | 15.7%     | 3.5%   | 22.1%  | 100.0%  |

## 1.4 WATER BODY DESIGNATED USE

Designated beneficial uses and water quality standards are established by the State of Mississippi in the *Water Quality Criteria for Intrastate, Interstate and Coastal Waters* regulations. These regulations set the criteria concentrations for pollutants and methods for calculating loads based on the standards. MDEQ regulations require the use of these standards for establishing loads for Mississippi waters. The standards for the listed segments of the Pascagoula River have been established based on a designated use of Fish and Wildlife.

## 1.5 APPLICABLE WATER QUALITY STANDARDS

Mercury is included within MDEQ regulations as a toxic substance. The standards specifically set the numeric criteria and calculation methods for determining the loading from sources for this pollutant.

The water quality standards applicable to the uses of the water body segment and the pollutant of concern are listed in Table 3 as defined by the current *State of Mississippi Water Quality Criteria for Intrastate, Interstate, and Coastal Waters* regulations.

Table 3. State of Mississippi Water Quality Criteria for Intrastate, Interstate, and Coastal Waters

| Parameter   | Beneficial use       | Water Quality Criteria   |
|---|----------------------|--|
| Total Mercury   | Public Water Supply  | Concentration may not exceed 0.151 µg/l  |
| Total Mercury   | Fish Consumption     | Concentration may not exceed 0.153 µg/l  |
| Mercury (II)<br>total dissolved Hg(II)<br>expressed as total<br>recoverable | Aquatic Life Support | <p><i>Fresh Water</i></p> <p><u>Acute</u>: instantaneous concentration may not exceed 2.1 µg/l</p> <p><u>Chronic</u>: average concentration may not exceed 0.012 µg/l expressed as total recoverable</p> <p><i>Salt Water</i></p> <p><u>Acute</u>: instantaneous concentration may not exceed 1.8 µg/l</p> <p><u>Chronic</u>: average concentration may not exceed 0.025 µg/l expressed as total recoverable</p> |

## 2.0 TMDL ENDPOINT AND WATER QUALITY ASSESSMENT

### 2.1 SELECTION OF A TMDL ENDPOINT AND CRITICAL CONDITION

One of the major components of a TMDL is the establishment of instream numeric endpoints, which are used to evaluate the attainment of acceptable water quality. Instream numeric endpoints, therefore, represent the water quality goals specified in the TMDL. The endpoints allow for a comparison between observed instream conditions and conditions that are needed to restore designated uses. However, due to the many unknowns within the mercury cycle, there is no clearly defined linkage between water column mercury loading and bioaccumulation rates within the fish. In the Executive Summary (Vol. I, Page O-2) of its Mercury Study report to Congress, EPA states that

*“given the current scientific understanding of the environmental fate and transport of this element, it is not possible to quantify how much of the methylmercury in fish consumed by the U.S. population is contributed by U.S. emissions relative to other sources of mercury (such as natural sources and re-emissions from the global pool). As a result, it cannot be assumed that a change in total mercury emissions will be linearly related to any resulting change in methylmercury in fish, nor over what time period these changes would occur.”<sup>12</sup>*

#### 2.1.1 Mercury Speciation and Chemistry

It has long been recognized that the chemical form of mercury (Hg) in air, water, and soil include elemental mercury Hg(0), inorganic ionic mercury (HgII) as soluble (HgIIs) or particulate mercury forms (HgIIp), and the organic form called monomethylmercury (MMHg or HgCh<sub>3</sub><sup>+</sup>). Each form has different behaviors that depend on its chemical and physical properties.<sup>4</sup>

The predominant source of mercury is atmospheric deposition. The atmospheric burden of mercury arises from both natural and anthropogenic sources accumulated over large periods. Both land and water environments release background mercury in the form Hg(0), except when combustion (forest and other terrestrial fires, fossil fuel combustion, waste combustion, etc.) produces the oxidized form – HgII. Hg(0) dissolves in water according to Henry’s Law, and is only weakly soluble in water (about 0.006 ng/l at equilibrium with present-day air concentrations).<sup>3</sup> Thus, Hg(0) must oxidize to HgII, which then is the predominant form of mercury in wet or dry deposition. Hg(0) has a half-life of about 1 year in the atmosphere, while that of HgII varies between hours to months.

Only a fraction of mercury entering watersheds from deposition actually is transported into waterbodies. Values ranging from 5 to 50 percent have been reported, and a common value of 25 percent has often been quoted.<sup>17</sup> Most of the mercury entering the watershed remains in the soil or terrestrial biota, or is reduced to Hg(0) and transfers back to the atmosphere by evasion. Thus, direct deposition on the water body frequently overshadows delivery from the watershed in many aquatic systems studied in the northern U.S.<sup>4</sup>

### **2.1.2 Mercury Transport and Transformations**

Mercury that makes its way into aquatic environments is essentially all inorganic ionic HgII. Hg(0) is only weakly soluble in water, while organic forms are usually present in trace amounts with MMHg in the typical range of 0.1 to 5 percent of the total mercury. However, higher amounts of MMHg can enter from wetland drainage.<sup>17</sup> Measurements of MMHg in rainwater seem to be associated with marine production of dimethylmercury, which hydrolyzes to form MMHg. Dimethylmercury does not seem to occur in fresh-water environments but only in the marine environment.<sup>5</sup> The ionized forms of mercury (HgII, MMHg) react rapidly and strongly with particulates. Furthermore, ionized forms react strongly with sulfide ions and somewhat strongly with organic complexes.

The production of MMHg by microorganisms and its subsequent accumulation in fish is by far the greatest concern. Part of that concern arises from MMHg's long biological half-lives in fish (1-2 years) as opposed to humans and other warm-blooded creatures that have half-lives of 1-3 months. Thus fish can accumulate MMHg to high levels, and the consumed fish – especially long-lived predatory fish – provide exposure of sensitive fish-eating organisms to MMHg.

Two competing processes affect the concentrations of MMHg, methylation produces MMHg while demethylation cleaves the methyl group and then reduces HgII to Hg(0) in a two-step process. The net MMHg produced is what scientists measure and organisms accumulate.

Microorganisms perform most of the methylation and demethylation, and sulfate reducing bacteria produce almost all of the MMHg.<sup>2</sup> The concentration of sulfate necessary to support production has an optimum because at higher concentrations, the produced sulfide binds HgII and can make it less available for uptake by sulfate reducing bacteria. Thus, many factors control the production of MMHg: the availability of HgII controlled largely by particulate material and dissolved organic carbon compounds; sulfide and sulfate concentrations; the presence of active sulfate reducing bacteria, and zones of sulfate production. MMHg production is often associated with sediments because most of the HgII is there and anaerobic conditions associated with reductive processes like sulfate reduction also occur there. The presence of sediments along with a ready source of biodegradable organic carbon resulting from plant production, may explain why wetlands are a major locale for production of MMHg. Circulation with surface waters may make wetland MMHg available for uptake. Emerging insects may substantially increase transfer of MMHg produced in wetlands to predatory fish.<sup>11</sup>

The food web has an important role in distributing MMHg into fish populations where fish consumers can then become part of the food web. The wide variability in mercury concentrations in similar sized fishes arise from the variety of local conditions of mercury bio-availability, MMHg production, and MMHg transfer among food web components.<sup>4</sup>

### 2.1.3 Mississippi Mercury Criteria (Fresh Water)

The beneficial use listed in Table 3 applicable to the listed segments of the Pascagoula River is Fish Consumption. The human health parameter for Fish Consumption is a total mercury concentration of 0.153 µg/l. The purpose of this standard is to restrict the mercury levels in fish tissue to below the 1.0 ppm FDA advisory level for human consumption. The total mercury human health standard of 0.153 µg/l in Mississippi's water quality standards was determined based on the accumulation of mercury in the types of fish that are commonly consumed in the state. Because the impaired segments are listed for partially supporting the use of Fish Consumption, the human health standard is an appropriate endpoint for Phase One of this TMDL study.<sup>8</sup>

However, the aquatic life criterion in fresh water, 0.012 µg/l of total Hg(II) is currently the more restrictive criterion for mercury concentration in the water column. We believe the toxicity criteria are overprotective of toxicity to aquatic life. According to *Ambient Water Quality Criteria for Mercury – 1984*, the 0.012 µg/l criterion for aquatic life was calculated based on a FDA action level of 1.0 mg/kg. This is a concentration of mercury in fish tissue of edible fish. The criterion was also based on a bioconcentration factor (BCF) of 81,700, which was the laboratory-determined ratio of the concentration of mercury in the tissue of the fathead minnow to the concentration of dissolved HgII in the lab water. The BCF of 81,700 is based on the transfer of mercury from the water to the tissue of the fathead minnow, and not directly to any species of edible fish.

In the “unused data” section of the same criteria document, BCF's ranging from 373 to 2400 were calculated for Bluegill, although the footnotes report that each BCF was not dependent on the concentration in the water. This means that there was no direct correlation between successive samples of mercury in the water and in the Bluegill fish tissue. However, a BCF was calculated in each case anyway, and they were much lower than the fathead minnow BCF. Although the criteria document states that the high BCF of the fathead minnow “might be more representative of commonly consumed warm-water fishes”; the Bluegill (which is a freshwater fish common in Mississippi) contradicts that assumption. To infer that the BCF of mercury in fathead minnows “might” be representative in light of the stated Bluegill results is an overprotective conclusion. Therefore, the use of the 0.012 µg/l of total mercury as the endpoint target for this TMDL incorporates an implicit margin of safety.

Additionally, we believe the 0.153 µg/l human health criterion is also protective of aquatic life. In *EPA's National Recommended Water Quality Criteria-Correction*, April 1999, EPA published 0.770 µg/l as the proposed freshwater aquatic life criterion. In effect, EPA has said that 0.153 µg/l is five times more protective of aquatic life than the proposed criterion. We believe 0.153 µg/l is protective of aquatic life while 0.012 µg/l is overprotective of aquatic toxicity, (a conclusion that EPA has supported by virtue of the latest proposed aquatic life criteria publication of 0.770 µg/l). If Mississippi's water quality criteria regarding mercury change, this Phase One TMDL will be revised to reflect those changes.



However, fish flesh sampling data indicate impairment of the water body's designated use. Therefore, to account for the uncertainty inherent with mercury fate and transport, this TMDL calls for a moratorium on future mercury discharges in the Pascagoula River watershed. This is to ensure the overall mercury load from point source contributors to the system does not increase. In addition, the TMDL includes an explicit MOS set at 50% for this TMDL.

## **2.2 MISSISSIPPI REGULATIONS ON FLOW DETERMINATION**

In addition to the endpoint, the flow rate must be determined in order to calculate the TMDL. According to Section II.9.D(2) of the *State of Mississippi Water Quality Criteria for Intrastate, Interstate and Coastal Waters* regulations, the 7Q10 flow shall be used when applying chronic toxicity criteria concentrations to calculations determining the load to a stream. However, portions of the Pascagoula River are tidally influenced. Statistically calculated flow values such as the 7Q10 flow are not valid under tidal influences.

The Mississippi Commission on Environmental Quality (MCEQ) has established a low-flow of 1,030 cfs for the Pascagoula River at Cumbest Bluff based on a correlation with data from USGS gaging station 02479000 (Pascagoula River at Merrill, MS). This is the most downstream point at which a 7Q10 flow is valid. As a conservative assumption, TMDL calculations made within this report will be based on this flow value.

## **2.3 DISCUSSION OF INSTREAM WATER QUALITY**

The listed segments of the Pascagoula River are listed based on fish tissue data collected from locations along the Pascagoula River, East Pascagoula River, and West Pascagoula River. Data collected at these sites are summarized and analyzed in the following sections.

### **2.3.1 Inventory of Water Quality Monitoring Data**

Fish tissue samples were collected by MDEQ from the following stations within the Pascagoula River, East Pascagoula River, and West Pascagoula River:

- Pascagoula River @ Road 614
- Pascagoula River @ Benndale
- Pascagoula River @ Cumbest Bluff
- Pascagoula River @ Poticaw Camp
- East Pascagoula River @ L.R. Marina
- West Pascagoula above I-10

Samples of various fish were collected between 1994 and 2001. These data are provided in Appendix A.

### 2.3.2 Analysis of Fish Tissue Data

Fish tissue data have been analyzed to identify violations requiring fish consumption advisories. Statistical summaries of methylmercury levels in fish tissue (wet weight filets) collected from each sampling location are presented in Table 4. These summaries are based on available data from 1994 to 2001, which is listed in Appendix A.

A single sampling event could have more than one fish, so the number of samples is listed along with the number of fish collected at that site. The percent exceedance value references the number of sampling events that averaged above the 1.0 ppm FDA action level. This percentage does not represent the number of individual fish that were found to exceed the action level. The table also gives the minimum, maximum, and average methylmercury levels found for all of the samples collected at the site. The fish tissue data collected from these water bodies are provided in Appendix A.

Table 4. Water Quality Station Data Analysis

| Station                             | Sample Events | Number of Fish | Percent Exceedance* | Min ppm | Max ppm | Average ppm |
|-------------------------------------|---------------|----------------|---------------------|---------|---------|-------------|
| Pascagoula River @ Road 614         | 6             | 8              | 17%                 | 0.48    | 1.13    | 0.71        |
| Pascagoula River @ Benndale         | 5             | 10             | 0%                  | 0.17    | 0.91    | 0.49        |
| Pascagoula River @ Cumbest Bluff    | 3             | 9              | 67%                 | 0.77    | 1.31    | 1.05        |
| Pascagoula River @ Poticaw Camp     | 19            | 38             | 42%                 | 0.45    | 1.73    | 1.00        |
| East Pascagoula River @ L.R. Marina | 6             | 17             | 17%                 | ND      | 1.31    | 0.65        |
| West Pascagoula River Above I-10    | 16            | 53             | 6%                  | ND      | 1.84    | 0.43        |

\* Percent exceedance is based on sampling events not individual fish.

### **3.0 SOURCE ASSESSMENT**

A TMDL evaluation must examine all known potential sources of the pollutant in the subject watershed, including point sources, nonpoint sources, and background levels. The source assessment is used as the basis of development of the model and ultimate analysis of the TMDL allocation options. However, in this Phase One Mercury TMDL, only point source contributions are considered for evaluation. Phase Two of the TMDL will further study contributions from nonpoint sources and background levels in the analysis.

#### **3.1 POTENTIAL SOURCES OF MERCURY**

Mercury emissions can occur from both natural and man-made sources. The man-made sources are estimated to account for the majority of all emissions. Appendix C contains a thorough outline of mercury sources. The following are examples of mercury sources in the environment that can be controlled.<sup>7</sup>

- Cement and Lime Kilns
- Coal and Oil Burning
- Copper Smelting
- Crematories
- Dental Amalgam Preparation/Disposal
- Dwelling Demolition (thermostats and switches)
- Electrical Product Manufacturing and Disposal (switches, fluorescent lights, some headlights and batteries)
- Evaporation of Mercury from Landfills
- Garbage Incinerators
- Hazardous Waste Incinerators
- Industrial Waste Discharge
- Laboratories Use and Waste
- Medical Waste Incinerators
- Petroleum Refining
- Residential Boilers
- Wastewater Treatment Plants and Sewage
- Wood Burning

Many items that we are in contact with everyday contain mercury. When these items are no longer useful, care should be taken to ensure that they are kept out of the trash or drain. When products containing mercury are placed in the trash, the mercury doesn't disappear. It finds its way into the environment from waste incinerators, landfills, or wastewater treatment facilities.

Items that may contain mercury include:

- Fluorescent Lamps
- Mercury Switches
- Mercury Vapor Lamps
- Thermostat Probes
- Metal Halide Lamps
- Relays
- High Pressure Sodium Lamps
- Thermometers
- Neon Lamps
- Thermostats
- Dental Amalgam
- Manometers
- Gauges
- Laboratory Solutions

### **3.1.1 Fluorescent and High-Intensity Discharge Lamps**

Fluorescent and high-intensity discharge (HID) lamps are used because they can use up to 50% less electricity than incandescent lighting. However, these lamps must be managed and disposed of properly because they contain mercury.<sup>8</sup>

### **3.1.2 Mercury Switches and Relays**

Mercury switches are found in a variety of items ranging from chest freezers to sump pumps. Mercury containing tilt switches are found under the lids of clothes washers and chest freezers. They stop the spin cycle or turn on a light. They are also found in motion-sensitive and position-sensitive safety switches in clothes irons and space heaters. Float switches are commonly used in sump pumps and bilge pumps to turn the equipment on and off when the water is at a certain level. Automobile trunk and hood light switches often contain mercury. A variety of manufacturing processes use relays to control power to heaters or pumps. Relays that contain mercury switches activate airbags, anti-lock brakes, some seat belt systems, and some automatically adjusting suspension systems. Some agricultural equipment, military vehicles, mass transit vehicles, and fire hook and ladder equipment also contain mercury switches.<sup>8</sup>

### **3.1.3 Mercury-Containing Thermostats and Thermostat Probes**

Mercury-containing tilt switches have been used in thermostats for more than 40 years. They provide accurate and reliable temperature control, require little or no maintenance, and do not require a power source. However, each switch contains approximately 3 grams of mercury. Mercury-free thermostats are available. Electronic thermostats now provide many of the same features as mercury thermostats.<sup>8</sup>

Mercury-containing thermostat probes may be found in several types of gas-fired appliances that have pilot lights, such as ranges, ovens, clothes dryers, water heaters, furnaces, or space heaters.

### 3.1.4 Mercury Thermometers

Some fever and laboratory thermometers contain mercury and should not be thrown in the trash. A typical fever thermometer contains about 0.5 grams of mercury. Larger laboratory thermometers can contain up to 3 grams of mercury. Many thermometers used to measure air and water temperature also contain mercury. They are used by homeowners, businesses, institutions, and recreational anglers. When the thermometers break outdoors, the mercury is difficult to capture. Mercury free thermometers such as digital thermometers are as accurate as mercury thermometers for most applications.<sup>8</sup>

### 3.1.5 Gauges, Manometers, Barometers, and Vacuum Gauges

Many barometers and vacuum gauges found in machinery contain mercury. Liquid mercury in the gauges responds to air pressure in a precise way that can be read on a calibrated scale. Several mercury-free alternatives are available. Some operate on the same principle as mercury gauges but use mercury-free liquids in the tube.<sup>8</sup>

Needle or bourdon gauges operate under a vacuum with a needle indicator. Electronic gauges can be used to measure pressure, but they must be calibrated with a mercury manometer. Equipment manufacturers recommend that service technicians use a needle or digital gauge to test the systems they are servicing, but that they calibrate the gauges they use in the field with a mercury manometer kept at their shop.<sup>8</sup>

## 3.2 POINT SOURCE ASSESSMENT

The point sources within the Pascagoula River watershed are listed in Table 5. Point sources that are possible contributors of mercury or that have flows greater than 0.05 MGD will be recommended by this TMDL to monitor their wastewater effluent for mercury. In an attempt to control mercury levels in the water body, this Phase One TMDL will call for a moratorium on any future increase in mercury discharges into the Pascagoula River watershed.

Table 5. Permitted Facilities in the Pascagoula River Watershed

| NPDES ID  | Facility Name  | County  | Receiving Water                   | Permitted Flow (MGD) |
|-----------|--|---------|-----------------------------------|----------------------|
| MS0031828 | George County Schools<br>Central Elementary School               | George  | Unnamed Tributary to<br>Big Creek | 0.0160               |
| MS0044504 | Lucedale POTW  | George  | Big Cedar Creek                   | 0.5000               |
| MS0020249 | MS Gulf Coast Regional Wastewater<br>Authority - Pascagoula POTW | Jackson | Pascagoula River                  | 10.0000              |

Table 5 Cont'd. Permitted Facilities in the Pascagoula River Watershed

| <b>NPDES ID</b> | <b>Facility Name</b>   | <b>County</b> | <b>Receiving Water</b>                             | <b>Permitted Flow (MGD)</b> |
|-----------------|--|---------------|--|-----------------------------|
| MS0028355       | Jackson County School District<br>East Central School          | Jackson       | Black Creek  | 0.0500                      |
| MS0028762       | Jackson County School District<br>Vancleave Junior High School | Jackson       | Unnamed Tributary to<br>Bluff Creek                | 0.0240                      |
| MS0032115       | Bluff Creek Mobile Home Park                                   | Jackson       | Bluff Creek  | 0.0340                      |
| MS0033375       | Kwik Kar Wash  | Jackson       | Unnamed Tributary to<br>Mary Walker Bayou          | 0.0009                      |
| MS0035637       | Gulf City Seafoods, Inc.                                       | Jackson       | Pascagoula River                                   | 0.0297*                     |
| MS0035769       | MDOT Interstate 10 Rest Area                                   | Jackson       | Marsh to Farrigut Lake to<br>West Pascagoula River | 0.0500                      |
| MS0038326       | Jackson County School District<br>Vancleave High School        | Jackson       | Unnamed Tributary to Bluff<br>Creek                | 0.0165                      |
| MS0042269       | Finicky Pet Food, Inc.   | Jackson       | East Pascagoula River                              | 0.0040*                     |
| MS0043010       | MS Gulf Coast Regional Wastewater<br>Authority - Gautier POTW  | Jackson       | West Pascagoula River                              | 4.0000                      |
| MS0043966       | Seachick, Inc.   | Jackson       | Unnamed Tributary to Clarke<br>Bayou               | 4.4300*                     |
| MS0046221       | Davis Laundry and Quickstop                                    | Jackson       | Black Creek  | 0.0036                      |
| MS0047007       | Circle V Traditions, Inc.                                      | Jackson       | Unnamed Tributary to Little<br>Bluff Creek         | 0.0024                      |
| MS0048445       | Colle Towing Company, Inc.                                     | Jackson       | Pascagoula River                                   | 0.0050                      |
| MS0055379       | Pascagoula Water Treatment Plant -<br>Communny Avenue          | Jackson       | Unnamed Creek to Yazoo Lake                        | 60.4500*                    |

\* This NPDES facility does not have a permit limit for flow. The facility is required to report flow information in each discharge monitoring report. The flow value given above is an average of the facility's historical flow data.

NPDES regulated construction activities and Municipal Separate Storm Sewer Systems (MS4s) are also considered point sources to surface waters and were included in this point source assessment. As of March 2003, discharge of storm water from construction activities disturbing between one and five acres must also be controlled by an NPDES permit. The purpose of these NPDES permits is to eliminate or minimize the discharge of pollutants from construction activities. Since construction activities at a site are of a temporary, relatively short term nature, the number of construction sites covered by the general permit varies. The NPDES regulated construction activities and MS4s will be regulated as specified in Mississippi's General Stormwater Permits for Small Construction, Construction, and Phase I & II MS4 permits.

There are four Phase II MS4s in the Pascagoula River watershed as well as one NPDES facility permitted to discharge stormwater into the Pascagoula River watershed. The discharges from these sources occur in response to storm events. Therefore, it is difficult to estimate the contribution from these sources.

The permitted stormwater sources in the Pascagoula River watershed are:

- Gautier MS4
- Jackson County MS4
- Moss Point MS4
- Pascagoula MS4
- Northrop Grumman Ship Systems, Inc. – NPDES Facility MS0003069

## **4.0 MODELING PROCEDURE**

Establishing the relationship between the instream water quality target and the source loadings is a critical component of TMDL development. It allows for the evaluation of alternatives for possible wasteload reductions. The link for mercury in the water column and mercury in fish flesh has not been established. The discussion of mercury TMDL calculations is included in this section.

### **4.1 MODELING CALCULATIONS**

Mass balance equations have been used to determine the mercury TMDL in the Pascagoula River watershed. A more complicated model is not warranted for Phase One of the TMDL analyzed because: (1) only contributions from point sources are considered, but none are known; (2) the mercury cycling processes will not be represented until Phase Two; (3) and water quality data for ambient mercury concentrations are not available to correspond to the levels of mercury found in the fish flesh for the Pascagoula River.

### **4.2 CALCULATION SETUP**

The watershed for the listed segments of the Pascagoula River contains all of HUC 03170006. The delineation of the watershed is based primarily on an analysis of the National Hydrography Dataset (NHD) stream network in the watershed. All upstream tributaries and point source loads located within HUC 03170006 are included in this TMDL.

As a conservative assumption, the 7Q10 low flow condition is the flow used to calculate this TMDL. However, portions of the Pascagoula River are tidally influenced. Statistically calculated flow values such as the 7Q10 flow are not valid under tidal influences. MDEQ has established a low-flow of 1,030 cfs for the Pascagoula River at Cumbest Bluff. This number is from a correlation with data from USGS gaging station 02479000 (Pascagoula River at Merrill, MS) and is the most downstream point at which a 7Q10 flow is valid. By using this conservative approach to calculating the flow, the implicit margin of safety is increased. TMDL calculations made within this report will be based on this flow value.

In addition, the MDEQ Office of Land and Water has taken flow measurements within the Pascagoula River, the West Pascagoula River, and the East Pascagoula in order to better characterize the division of flow. These measurements indicated that the flow in the Pascagoula River is split with approximately 60% flowing to the West Pascagoula River and the remaining 40% flowing to the East Pascagoula River.

### **4.3 SOURCE REPRESENTATION**

Only point sources are considered in this Phase One Mercury TMDL. Point sources that are possible contributors of mercury or that have flows greater than 0.05 MGD will be recommended by this TMDL to monitor their wastewater effluent for mercury. Table 6 lists the facilities that are recommended for mercury monitoring in the Pascagoula River watershed.



Table 6. Facilities in the Pascagoula River Watershed Recommended for Mercury Monitoring

| <b>NPDES ID</b> | <b>Facility Name</b>  |
|-----------------|---|
| MS0031828       | George County Schools - Central Elementary School             |
| MS0044504       | Lucedale POTW   |
| MS0020249       | MS Gulf Coast Regional Wastewater Authority - Pascagoula POTW |
| MS0028355       | Jackson County School District - East Central School          |
| MS0028762       | Jackson County School District - Vancleave Junior High School |
| MS0035769       | MDOT Interstate 10 Rest Area                                  |
| MS0038326       | Jackson County School District - Vancleave High School        |
| MS0043010       | MS Gulf Coast Regional Wastewater Authority - Gautier POTW    |
| MS0055379       | Pascagoula Water Treatment Plant - Communny Avenue            |

A significant amount of mercury water quality sampling data from the Pascagoula River is needed to adequately explain the relationship between the mercury concentration in the water column with the concentration in fish tissue. As ambient mercury data and tools for analyzing mercury cycling become available, Phase Two of this TMDL project will be completed to accurately represent mercury sources, atmospheric deposition, and stream response.

## 5.0 ALLOCATION

TMDLs are composed of the sum of individual waste load allocations ( $\Sigma$ WLA) for point sources, the sum of load allocations ( $\Sigma$ LA) for nonpoint sources, and a margin of safety (MOS). This definition is mathematically expressed by the equation:

$$\text{TMDL} = \Sigma\text{WLA} + \Sigma\text{LA} + \text{MOS}$$

The TMDL is the amount of a pollutant that can be assimilated by the receiving water while still achieving water quality standards. This TMDL represents the maximum load of mercury that can be introduced into the water body by point source discharge based on Mississippi's mercury criterion.

### 5.1 TMDL CALCULATION

The TMDL Calculation is based upon the conservation of mass principle, where the load can be calculated by using the following relationship:

$$\text{Concentration} = \text{Load} / \text{Flow}$$

Rearranging this equation, the load can be calculated as follows:

$$\text{Load} = \text{Concentration} * \text{Flow}$$

$$\text{Load gm/day} = 0.012 \mu\text{g/l} * 1,030 \text{ cfs} * 2.45 \text{ (unit conversion factor)} = 30.3 \text{ gm/day}$$

The overall TMDL load for total mercury in the water body system is 30.3 grams per day. The total mercury II target of 0.012  $\mu\text{g/l}$  is expressed as Total Recoverable Mercury. As mentioned earlier, the MDEQ Office of Land and Water has taken flow measurements within the Pascagoula River, the West Pascagoula River, and the East Pascagoula in order to better characterize the division of flow. These measurements indicated that the flow in the Pascagoula River is split with approximately 60% flowing to the West Pascagoula River and the remaining 40% flowing to the East Pascagoula River.

Table 7. TMDL for Total Mercury II

| Water Body            | Segment ID             | Flow (cfs) | Total Hg(II) Target ( $\mu\text{g/l}$ ) | TMDL (gm/day) |
|-----------------------|------------------------|------------|---|---------------|
| Pascagoula River      | MSPASRM1               | 1030       | 0.012                                   | 30.3          |
| East Pascagoula River | MSEPASRM1<br>MSEPASRM2 | 412        | 0.012                                   | 12.1          |
| West Pascagoula River | MSWPASRM1<br>MSWPASRM2 | 618        | 0.012                                   | 18.2          |

## **5.2 TMDL ALLOCATIONS**

The allocation for this TMDL involves a wasteload allocation for point sources and a load allocation for nonpoint sources necessary for attainment of water quality standards in the Pascagoula River watershed. The Wasteload Allocation for this TMDL was determined by multiplying the permitted flow of the dischargers within the Pascagoula River watershed by the mercury criterion. Since this TMDL calls for a moratorium on any future increase in mercury discharges into the Pascagoula River watershed, the WLA component should not increase. The Load Allocation accounts for atmospheric deposition and background.

### **5.2.1 Wasteload Allocations**

The sum of the loads allocated to the point sources, ( $\Sigma$ WLAs) is determined by multiplying the permitted flow from the facility by the mercury criterion. Appendix B provides a list of permitted facilities within the watershed. Each facility's flow is multiplied by the mercury criterion and a conversion factor to give a daily load. This TMDL recommends that possible contributors of mercury or facilities with flows greater than 0.05 MGD monitor wastewater effluent for mercury.

All municipalities must complete the NPDES Form 2A Application. According to this application, supplemental information regarding effluent testing and toxicity testing must be included if one or more of the following is true: (1) the facility has a design flow rate greater than or equal to 1.0 MGD, (2) the facility is required to have a pretreatment program, or (3) the facility is otherwise required by the permitting authority to provide additional effluent information or submit the results of toxicity testing. Some facilities listed below will be required to monitor for mercury in order to meet their 2A application requirements. Minor municipal facilities not meeting any of the above requirements are not required to complete all of the 2A application are not required to monitor for mercury. In addition, the 2A application does not apply to commercial and industrial facilities. However, due to the mercury impairment in this watershed, this TMDL recommends that mercury monitoring similar to that required by a complete 2A application be required for all facilities listed in Table 8. This TMDL recommends that all facilities listed below have mercury monitoring required as part of their permitting process. If mercury is found in the facility's discharge, permit limits could be developed and a mercury minimization program would be needed.

Table 8. Facilities Recommended to Perform Mercury Effluent Monitoring

| Facility Name | NPDES ID  |
|---------------|---|
| MS0031828     | George County Schools - Central Elementary School             |
| MS0044504     | Lucedale POTW   |
| MS0020249     | MS Gulf Coast Regional Wastewater Authority - Pascagoula POTW |
| MS0028355     | Jackson County School District - East Central School          |
| MS0028762     | Jackson County School District - Vancleave Junior High School |
| MS0035769     | MDOT Interstate 10 Rest Area                                  |
| MS0038326     | Jackson County School District - Vancleave High School        |
| MS0043010     | MS Gulf Coast Regional Wastewater Authority - Gautier POTW    |
| MS0055379     | Pascagoula Water Treatment Plant - Communy Avenue             |

NPDES regulated construction activities and MS4s are also considered point sources to surface waters and were included in the WLA component of this TMDL. As of March 2003, discharge of storm water from construction activities disturbing between one and five acres must also be controlled by an NPDES permit. The purpose of these NPDES permits is to eliminate or minimize the discharge of pollutants from construction activities. Since construction activities at a site are of a temporary, relatively short term nature, the number of construction sites covered by the general permit varies. The NPDES regulated construction activities and MS4s will be regulated as specified in Mississippi’s General Stormwater Permits for Small Construction, Construction, and Phase I & II MS4 permits.

There are four Phase II MS4s in the Pascagoula River watershed as well as one NPDES facility permitted to discharge stormwater into the Pascagoula River watershed. The discharges from these sources occur in response to storm events. Therefore, it is difficult to estimate the contribution from these sources.

The permitted stormwater sources in the Pascagoula River watershed are:

- Gautier MS4
- Jackson County MS4
- Moss Point MS4
- Pascagoula MS4
- Northrop Grumman Ship Systems, Inc. – NPDES Facility MS0003069

### **5.2.2 Load Allocations**

The Load Allocations (LA) for this TMDL account for mercury due to atmospheric deposition and background. Since atmospheric deposition is believed to be the primary source of mercury, a large portion of the TMDL has been set aside for this component. Phase Two of this TMDL project will explore atmospheric deposition along with local and national air-emission reduction goals.

### 5.3 INCORPORATION OF A MARGIN OF SAFETY

The two options for MOS development are either to implicitly incorporate the MOS using conservative assumptions or to explicitly specify a portion of the total TMDL as the MOS. A dual MOS method has been selected for this Phase One TMDL. It is implicit, based on the conservative assumptions inherent in the selection of the TMDL endpoint of 0.012 µg/l. Calculating the TMDL based on the 7Q10 flow at the most downstream location available along the Pascagoula River is also a conservative assumption. This flow occurs above the Pascagoula River split and does not include flow contributions from tributaries entering below this point. In addition, this 7Q10 value does not account for additional flows entering the river from Pascagoula Bay due to tidal influences. As noted, this TMDL uses the 7Q10 flow and the allowable mercury concentration to determine the allowable load. Therefore, use of this conservative flow value results in a conservative load being assigned to these water bodies.

In addition, an explicit MOS has been included to account for uncertainty in the mercury linkage between fish flesh mercury levels and water-column mercury levels. The explicit MOS has been set at 50%.

Additional conservative assumptions for TMDL calculation are inherent in the development of the 0.153 µg/l human health standard. The criterion is based on the following equation:

$$C = \frac{\text{reference dose} * \text{human body weight}}{\text{fish consumption rate} * \text{bio-concentration factor}}$$

The criterion was based on a combination of fish consumption rates and bio-concentration factors for fresh water fishes, coastal organisms, and salt-water fishes. If the coastal organisms and salt-water fishes are omitted from the calculation, the criterion would be 2.22 µg/l. The fish tissue data collected from the Pascagoula River watershed, Mississippi show elevated mercury levels in largemouth bass and flathead catfish. However, the BCF used in the criteria development considers four species of freshwater fish resulting in an average BCF of 5500, which is higher than that of either the bass or the catfish. Using the higher combined value in the denominator of the above equation, another conservative assumption is introduced into the calculations.

Additionally, the fresh water fish consumption rate established in the *Ambient Water Quality Criteria for Mercury* is 1.72 gm/day per person. Our regulations, however, require the use of 6.5 gm/day per person. This calculation would set the criterion at 0.587 µg/l as compared to the 0.153 µg/l in Mississippi's water quality standards. The use of a fish consumption rate of almost 3.8 times that for freshwater species alone introduces yet another conservative assumption which is already a part of the current human health standard for Mississippi.

However, there is enough uncertainty inherent to this entire process to justify the inclusion of an explicit MOS. As previously mentioned, this explicit MOS has been set at 50%.

$$\text{TMDL} = \text{WLA} + \text{LA} + \text{MOS}$$

$$30.3 \text{ gm/day} = 3.62 \text{ gm/day WLA} + 11.53 \text{ gm/day LA} + 15.15 \text{ gm/day MOS}$$

## **5.4 SEASONALITY**

Wet deposition is greatest in the winter and spring seasons. Mercury will be expected to fluctuate based on the amount and distribution of rainfall, and variability of localized and distant atmospheric sources. While a maximum daily load is established in this TMDL, the average annual load is of greatest significance since mercury bioaccumulation and the resulting risk to human health that results from mercury consumption is a long term phenomenon. Thus, daily or weekly inputs are less meaningful than total annual loads over many years. The use of annual load allows for integration of short-term or seasonal variability. Inputs will continue to be estimated through monitoring and modeling.

Methylation of mercury is expected to be highest during the summer. High temperatures and static conditions result in hypoxic and/or conditions that promote methylation. Based on this enhanced methylation and high predator feeding activity during the summer, mercury bioaccumulation is expected to be greatest during the summer. However, based on the refractory nature of mercury, seasonal changes in body burden would be expected to be slight. Inherent variability of mercury concentrations between individual fish of the same and/or different size categories is expected to be greater than seasonal variability.<sup>15</sup>

## **5.5 IMPLEMENTATION PLAN**

Implementation of this Phase One Mercury TMDL will differ from other types of TMDLs since atmospheric deposition is believed to be the primary pollutant source. This will involve MDEQ working with stakeholders to identify the most appropriate mechanisms to implement this TMDL project. MDEQ will cooperate with EPA concerning national initiatives and strategies, which will be important to implement regulatory controls on a national and international basis. Much monitoring, research, and regulation is in progress on the national level. MDEQ will consider these ongoing activities in implementing this and future phases of this TMDL project.

The ultimate reduction of mercury in the environment will take numerous years and is in line with the Bi-national Toxics Strategy, which sets a national challenge of 50% reduction of mercury releases to the air by 2006. Phase Two of this TMDL project will explore atmospheric deposition along with local and national air-emission reduction goals. Long-term monitoring of wet deposition rates and fish tissue in the water body segment will serve as environmental indicators to evaluate the effectiveness of the TMDLs and other parallel control measures.<sup>15</sup>

MDEQ also supports and encourages Pollution Prevention activities (P2 activities) as part of this implementation plan. P2 activities help alleviate costs and resources associated with controlling, removing, and managing mercury contamination in the environment. These activities include: (1) separating mercury-containing waste from the trash and save it for local household hazardous waste collection days, (2) taking mercury-containing items such as thermometers to a household hazardous waste collection facility, (3) removing mercury-containing items from households and schools (including student laboratories), and (4) conserving electricity (burning less coal and oil, which naturally contains mercury, for electricity will emit less mercury into the environment). Table 9 gives some examples of possible P2 alternatives for products containing mercury.

Table 9. Pollution Prevention (P2) Alternatives for Products Containing Mercury

| <b>Discards Known to Contain Mercury</b>  | <b>P2 Alternatives</b>   |
|---|--|
| Thermometers  | Red Bulb (Alcohol) Thermometers<br>Digital Thermometers  |
| Thermostats<br>(non-electric models)  | Electric Models  |
| Batteries<br>(old alkaline type prior to 1996)                                      | Recharge Alkaline Batteries<br>Mercury Free Batteries  |
| Button Batteries  | Mercury Free Button Batteries (Zinc air type)  |
| Silver Amalgam Waste  | Ask Your Dentist   |
| Quicksilver Maze Toy  | Mercury-Free Toys  |
| Old Latex Paints<br>(since 1990, mercury has been banned in latex paints)           | New Latex Paint  |
| Some Shoes that Light Up<br>(L.A. Gear's My Lil' Lights if bought before June 1994) | Mercury-Free Shoes   |
| Switches<br>(some light and appliance switches)                                     | Mechanical or Electrical Switches  |
| Contact Lens Solution Containing Thimerosal   | Mercury-Free Solution  |
| Lights<br>(fluorescent, high intensity discharge, and mercury vapor lamps)          | Energy Efficient Fluorescent Lights<br>(These lights still contain mercury. However, energy will be conserved thereby reducing mercury emissions from coal and oil combustion) |

## **6.0 CONCLUSION**

MDEQ will not approve any NPDES Permit application for the Pascagoula River drainage area that does not comply with the moratorium for additional mercury discharges into this segment. In addition, this TMDL recommends all dischargers that are possible contributors of mercury or that have flows greater than 0.05 MGD to monitor for mercury using clean techniques and accurate testing methods. This TMDL also recommends and encourages Pollution Prevention Alternatives/Activities that address possible sources of mercury within the Pascagoula River watershed.

Phase Two of this TMDL will include nonpoint sources of mercury, atmospheric deposition, and will consider the effects of mercury cycling in the water body. The TMDL calculations from Phase One may be revised in Phase Two of this TMDL since more will be known about the percentage of mercury contributions from point and nonpoint sources.

### **6.1 FOLLOW-UP MONITORING**

Additional ambient mercury monitoring for all species of mercury will be needed for development of Phase Two. Additional information is required to facilitate the understanding of the methylmercury process and the linkage between mercury in the water column and mercury in fish flesh. Specialized monitoring approaches will also be needed to determine the atmospheric deposition contribution to mercury in the watershed.

MDEQ has adopted the Basin Approach to Water Quality Management, a plan that divides Mississippi's major drainage basins into five groups. During each yearlong cycle, MDEQ resources for water quality monitoring are focused on one of the basin groups. During the next monitoring phase in the Pascagoula River Basin, the Pascagoula River may receive additional monitoring to identify the improvements in water quality gained from the implementation of the Phase One strategy included in this TMDL. MDEQ plans to continue monitoring mercury concentrations in fish tissue within these segments of the Pascagoula River, East Pascagoula River, and West Pascagoula River in order to protect human health.

### **6.2 PUBLIC PARTICIPATION**

This Phase One TMDL project will be published for a 30-day public notice. During this time, the public will be notified by publication in both a statewide and local newspaper. The public will be given an opportunity to review the TMDL and submit comments. MDEQ also distributes all TMDLs at the beginning of the public notice to those members of the public who have requested to be included on a TMDL mailing list. TMDL mailing list members may request to receive the TMDL reports through either, email or the postal service. Anyone wishing to become a member of the TMDL mailing list should contact Greg Jackson at (601) 961-5098 or [Greg\\_Jackson@deq.state.ms.us](mailto:Greg_Jackson@deq.state.ms.us).

All comments received during the public notice period and at any public hearings become a part of the record of this TMDL. All comments will be considered in the ultimate approval of this TMDL and for submission of this TMDL to EPA Region 4 for final approval.



## DEFINITIONS

**Ambient Stations:** Network of fixed monitoring stations established for systematic water quality sampling at regular intervals, and for uniform parametric coverage over a long-term period.

**Assimilative Capacity:** The amount of contaminant load that can be discharged to a specific stream or river without violating the provisions of the *State of Mississippi Water Quality Criteria for Intrastate, Interstate, and Coastal Waters and Water Quality* regulations. Assimilative capacity is the extent to which a body of water can receive wastes without significant deterioration of beneficial uses.

**Atmospheric Deposition:** Input of chemical components from the atmosphere into natural waters through the processes of wet deposition (rain, snow) and dry deposition (particle fallout, gas-water exchange). Components can include nutrients, acidity, trace elements, and anthropogenic organics.

**Background:** Ambient pollutant concentrations due to natural sources, nearby sources other than the one currently under consideration, and unidentified anthropogenic sources.

**Best Management Practices:** (1) The methods, measures, or practices selected by an agency to meet its nonpoint source control needs. BMPs include but are not limited to structural and nonstructural controls and operation and maintenance procedures. BMPs can be applied before, during, or after pollution-producing activities to reduce or eliminate the introduction of pollutants into receiving waters. (2) Methods have been determined to be the most effective, practical means of preventing or reducing pollution from nonpoint sources.

**Bioaccumulation (Bioretention):** (1) The uptake and, at least temporary, storage of a chemical by an exposed animal. The chemical can be retained in its original form and/or as modified by enzymatic and non-enzymatic reactions in the body. (2) The retention and concentration of a substance by an organism. Accumulation of a large amount of a substance in the body by ingesting small amounts of the substance over an extended time. (3) The process by which a compound is taken up by an aquatic organism, both from water and through food.

**Bioaccumulation Factor (BAF):** The ratio of a substance's concentration in tissue versus its concentration in ambient water, in situations where the organism and the food chain are exposed.

**Bioconcentration:** (1) The accumulation of a chemical in tissues of an organism (such as fish) to levels greater than in the surrounding medium (such as water) in which the organism resides. (2) The net accumulation of a substance directly from water in to and onto aquatic organisms. (3) The process by which a compound is absorbed from water through the gills or epithelial tissues and is concentrated in the body.

**Bioconcentration Factor (BCF):** (1) The ratio of the concentration of a chemical in aquatic organisms to the amount in water at equilibrium. (2) The measure of the tendency for a substance to accumulate in the tissue of an aquatic organism. BCF is determined by the extent of partitioning of a substance, at equilibrium, between the tissue of an aquatic organism and water. As the ratio of concentration of a substance in the organism divided by the concentration in water, higher BCF values reflect a tendency for substances to accumulate in the tissue of aquatic organisms.

**Calibration:** Testing and tuning of a model to a set of field data. Also includes minimization of deviations between measured field conditions and output of a model by selecting appropriate model coefficients.

**Critical Condition:** Hydrologic and atmospheric conditions in which the pollutants causing impairment of a water body have their greatest potential for adverse effects.

**Daily Discharge:** The "discharge of a pollutant" measured during a calendar day or any 24-hour period that reasonably represents the calendar day for purposes of sampling. For pollutants with limitations expressed in units of mass, the "daily discharge" is calculated as the total mass of the pollutant discharged over the day. For pollutants with limitations expressed in other units of measurement, the "daily average" is calculated as the average.

**Designated Use:** (1) Those uses specified in water quality standards for each water body or segment whether or not they are being attained. (2) Those water uses identified in state water quality standards which must be achieved and maintained as required under the Clean Water Act.

**Discharge Monitoring Report:** Report of effluent characteristics submitted by a facility that has been granted an NPDES Permit.

**Effluent:** (1) Any solid, liquid, or gas which enters the environment as a by-product of a man-oriented process. The substances that flow out of a designated source. Effluent, effluence, and efflux have the same meaning. (2) Wastewater – treated or untreated – that flows out of a treatment plant, sewer, or industrial outfall. Generally refers to wastes discharged into surface waters.

**Effluent Standards and Limitations:** All State or Federal effluent standards and limitations on quantities, rates, and concentrations of chemical, physical, biological, and other constituents to which a waste or wastewater discharge may be subject under the Federal Act or the State law. This includes, but is not limited to, effluent limitations, standards of performance, toxic effluent standards and prohibitions, pretreatment standards, and schedules of compliance.

**Geometric Mean:** The  $n$ th root of the product of  $n$  numbers. A 30-day geometric mean is the 30<sup>th</sup> root of the product of 30 numbers.

**Impairment:** Conditions in which the applicable state water quality standards are not met for a water body and the designated use is impaired.

**Load Allocation (LA):** The portion of a receiving water's loading capacity attributed either to one of its existing or future nonpoint sources of pollution or to natural background sources. Load allocations are best estimates of the loading, which may range from reasonably accurate estimates to gross allotments, depending on the availability of data and appropriate techniques for predicting the loading. Wherever possible, natural and nonpoint source loads should be distinguished.

**Loading:** (1) An amount of matter or thermal energy that is introduced into a receiving water. Loading may be either man caused (pollutant loading) or natural (background loading). (2) The concentration of a pollutant, usually expressed in grains of pollutant per cubic foot of contaminated gas stream.

**Margin of Safety (MOS):** A required component of the TMDL that accounts for the uncertainty about the relationship between the pollutant load and the quality of the receiving water body.

**Mercury (Hg):** A silver-white metal, atomic weight 200.59, which is a slightly volatile liquid at room temperature. Mercury is a naturally occurring element that is found in air, water and soil. It ranks about 67th in natural abundance among the elements in crustal rocks. Most of the mercury in the atmosphere is elemental mercury vapor (which circulates in the atmosphere for up to a year, and hence can be widely dispersed and transported thousands of miles from likely sources of emission). Most of the mercury in water, soil, sediments, or plants and animals is in the form of inorganic water-soluble salts (most commonly mercuric chloride) and organic forms of mercury (commonly methylmercury). Among the commercially important compounds of mercury are mercuric sulfide, a common antiseptic also used as the pigment vermilion; mercurous chloride, or calomel, used for electrodes, and formerly used as a cathartic; mercuric chloride, or corrosive sublimate; and medicinals such as Mercurochrome.

**Mercury (elemental):** Mercury in a zero (0) oxidation state - referred to as mercury vapor when present in the atmosphere and as metallic mercury when present in its liquid form.

**Mercury II (inorganic mercury):** Mercury which has been naturally oxidized to a divalent oxidation state and exhibits a wide range of acute toxicity to aquatic life. Inorganic mercury occurs in numerous forms/compounds; the most common include mercuric chloride ( $\text{HgCl}_2$ ), mercurous chloride ( $\text{Hg}_2\text{Cl}_2$ ), and mercuric oxide ( $\text{Hg}[\text{O}]$ ).

**Methylmercury (organic mercury):** Mercury II which has been methylated in surface waters by naturally occurring bacteria and which can substantially accumulate in the food chain. Nearly all of the mercury that accumulates in fish tissue is methylmercury.

**Nonpoint Source Pollution:** The pollution from sources which generally are not controlled by establishing effluent limitations under sections 301, 302, and 402. Nonpoint source pollutants are not traceable to a discrete identifiable origin, but generally result from land runoff, precipitation, drainage, or seepage. This water may contain pollutants that come from land use activities such as agriculture, construction, silviculture, surface mining, disposal of wastewater, hydrologic modifications, and urban development.

**NPDES Permit:** An individual or general permit issued by the MDEQ Permit Board pursuant to regulations adopted by the Commission under Mississippi Code Annotated (as amended) § 49-17-17 and § 49-17-29 for discharges into State waters.

**Part Per Million:** One millionth of a measurement. This nomenclature also applies to part per billion and part per trillion. 1 mg/kg mercury in fish flesh is one part per million. 1 µg/l liquid concentration is equivalent to one part per billion. 1 nanogram liquid concentration is equivalent to one part per trillion.

**Phased TMDL Project:** Under the phased approach, the TMDL has load allocations and wasteload allocations calculated with margins of safety to meet water quality standards. The allocations are based on estimates that use available data and information, but monitoring for collection of new data is required. The phased approach provides for further pollution reduction without waiting for new data collection and analysis.

**Point Source Pollution:** Pollution from a stationary location or fixed facility from which pollutants are discharged or emitted. Pollution from any single identifiable source, e.g., a pipe, ditch, ship, ore pit, or factory smokestack.

**Pollution Prevention (P2) Activities:** (1) The use of materials, processes, or practices that reduce or eliminate the creation of pollutants or wastes. It includes practices that reduce the use of hazardous materials, energy, water, or other resources, and practices that protect natural resources through conservation or more efficient uses. (2) A preventative action or a measure taken to minimize waste generation or waste toxicity, if waste generation is inevitable. In a broad sense, it incorporates clean technology, low-waste technology, non-waste technology, prevention, quantity reduction, recycling, reduction, resource conservation, resource recovery, source reduction, toxicity reduction, waste minimization, etc.

**Pollution:** Generally, the presence of matter or energy whose nature, location, or quantity produces undesired environmental effects. Under the Clean Water Act, for example, the term is defined as man-made or man-induced alteration of the physical, biological, and radiological integrity of water. Other pollution related terms include: agricultural pollution, air pollution, indoor air pollution, industrial waste pollution, manmade air pollution, natural pollution, noise pollution, oil pollution, sewage pollution, soil pollution, thermal pollution, water pollution, and wood burning stove pollution.

**Practical Bioaccumulation Factor (PBCF):** A practical approximation used in lieu of a BCF in the derivation of the human health criteria for mercury in Ambient Water Quality Criteria for Mercury. The PBCF's were calculated as the ratio of the average concentration of mercury in muscle in one species of fish to the average concentration of mercury in the body of water in which the species normally lives.

**Publicly Owned Treatment Works (POTW):** The treatment works treating domestic sewage that is owned by a municipality or State.

**Scientific Notation (exponential notation):** Mathematical method in which very large numbers or very small numbers are expressed in a more concise form. The notation is based on powers of ten. Numbers in scientific notation are expressed as the following:  $4.16 \times 10^{(+b)}$  and  $4.16 \times 10^{(-b)}$  [same as 4.16E4 or 4.16E-4]. In this case,  $b$  is always a positive, real number. The  $10^{(+b)}$  tells us that the decimal point is  $b$  places to the right of where it is shown. The  $10^{(-b)}$  tells us that the decimal point is  $b$  places to the left of where it is shown. For example:  $2.7 \times 10^4 = 2.7E+4 = 27000$  and  $2.7 \times 10^{-4} = 2.7E-4 = 0.00027$ .

**Sigma ( $\Sigma$ ):** Shorthand way to express taking the sum of a series of numbers. For example, the sum or total of three amounts 24, 123, 16, ( $d_1$ ,  $d_2$ ,  $d_3$ ) respectively could be shown as:

$$\sum_{i=1}^3 d_i = d_1 + d_2 + d_3 = 24 + 123 + 16 = 163$$

**STORET:** EPA's computerized water quality data base that includes physical, chemical, and biological data measured in water bodies throughout the United States.

**Storm Water:** (1) The storm water runoff, snow melt runoff, and surface runoff and drainage. (2) Rainfall that does not evaporate or infiltrate the ground because of impervious land surfaces or a soil infiltration rate than rainfall intensity, but instead flows into adjacent land or water bodies or is routed into a drain or sewer system.

**Total Maximum Daily Load (TMDL):** (1) The total allowable pollutant load to a receiving water such that any additional loading will produce a violation of water quality standards. (2) The sum of the individual waste load allocations and load allocations. A margin of safety is included with the two types of allocations so that any additional loading, regardless of source, would not produce a violation of water quality standards.

**Waste:** Useless, unwanted, or discarded material resulting from (agricultural, commercial, community, and industrial) activities. Wastes include solids, liquids, and gases.

**Wasteload Allocation (WLA):** (1) The portion of a receiving water's loading capacity that is allocated to one of its existing or future point sources of pollution. WLAs constitute a type of water quality based effluent limitation. (2) The portion of a receiving water's total maximum daily load that is allocated to one of its existing or future point sources of pollution. (3) The maximum load of pollutants each discharger of waste is allowed to release into a particular waterway. Discharge limits are usually required for each specific water quality criterion being, or expected to be, violated. The portion of a stream's total assimilative capacity assigned to an individual discharge.

**Water Quality Criteria:** Specific levels of water quality which, if reached, are expected to render a body of water suitable for its designated use. The criteria are based on specific levels of pollutants that would make the water harmful if used for drinking, swimming, farming, fish production, or industrial processes. Water quality criteria are comprised of numeric and narrative criteria. Numeric criteria are scientifically derived ambient concentrations developed by EPA or States for various pollutants of concern to protect human health and aquatic life. Narrative criteria are statements that describe the desired water quality goal.

**Water Quality Standards:** (1) Provisions of State or Federal law which consist of a designated use or uses for the water quality criteria for such waters based upon such uses. Water quality standards are to protect the public health or welfare, enhance the quality of water, and serve the purposes of the Clean Water Act. (2) A law or regulation that consists of the beneficial designated use or uses of a water body, the numeric and narrative water quality criteria that are necessary to protect the use or uses of that particular water body, and an antidegradation statement. (3) State-adopted and EPA-approved ambient standards for water bodies. The standards prescribe the use of the water body and establish the water quality criteria that must be met to protect designated uses.

**Waters of the State:** All waters within the jurisdiction of this State, including all streams, lakes, ponds, wetlands, impounding reservoirs, marshes, watercourses, waterways, wells, springs, irrigation systems, drainage systems, and all other bodies or accumulations of water, surface and underground, natural or artificial, situated wholly or partly within or bordering upon the State, and such coastal waters as are within the jurisdiction of the State, except lakes, ponds, or other surface waters which are wholly landlocked and privately owned, and which are not regulated under the Federal Clean Water Act (33 U.S.C.1251 et seq.).

**Watershed:** (1) The land area that drains (contributes runoff) into a stream. (2) The land area that drains into a stream; the watershed for a major river may encompass a number of smaller watersheds that ultimately combine at a common delivery point.

## **ABBREVIATIONS**

|        |   |
|--------|---|
| 7Q10   | Seven-Day Average Low Stream Flow with a Ten-Year Occurrence Period |
| BASINS | Better Assessment Science Integrating Point and Nonpoint Sources    |
| BMP    | Best Management Practice  |
| DMR    | Discharge Monitoring Report   |
| EPA    | Environmental Protection Agency                                     |
| GIS    | Geographic Information System                                       |
| HCR    | Hydrograph Controlled Release Facility                              |
| HUC    | Hydrologic Unit Code  |
| LA     | Load Allocation   |
| MARIS  | State of Mississippi Automated Resource Information System          |
| MDEQ   | Mississippi Department of Environmental Quality                     |
| MOS    | Margin of Safety  |
| MS4    | Municipal Separate Storm Sewer Systems                              |
| NHD    | National Hydrography Dataset  |
| NRCS   | National Resource Conservation Service                              |
| NPDES  | National Pollution Discharge Elimination System                     |
| NPSM   | Nonpoint Source Model   |
| P2     | Pollution Prevention  |
| PCS    | Permit Compliance System  |
| PPB    | Part per Billion ( $1 \times 10^{-9}$ ) ( $\mu\text{g/l}$ )         |
| PPM    | Part per Million ( $1 \times 10^{-6}$ ) ( $\text{mg/l}$ )           |
| PPT    | Part per Trillion ( $1 \times 10^{-12}$ ) ( $\text{ng/l}$ )         |
| USGS   | United States Geological Survey                                     |
| WLA    | Waste Load Allocation   |

## REFERENCES

1. ATSDR. 1999. Toxicological profile for mercury (update). Agency for Toxic Substances and Disease Registry, 1999
2. Gilmour, C.C., E. A. Henry, and R. Mitchell. 1992. Sulfate Stimulation of Mercury Methylation in Freshwater Sediments. *Environ. Sci. Technol.* 26:2281-2287.
3. Hudson, r. J. M., S. A. Gherini, c. J. Watras, et al. 1994. Modeling the biogeochemical cycle of mercury in lakes: The mercury cycling model (MCM) and its application to the MTL study lakes. *Mercury as a Global Pollutant*. Watras C. J. and J. W. Juckabee (Eds.). Lewis Publishers. Pp 473-523.
4. J. David Dean, Kent Thornton, Randy Manning, & Donald Porcella, "Mercury Issues in the Southeastern United States." 2000, a report prepared for the TAPPI meeting.
5. Mason, R. P. and W. F. Fitzgerald. 1990. Alkylmercury species in the equatorial Pacific. *Nature*. 347:457-459.
6. Metcalf and Eddy. 1991. *Wastewater Engineering: Treatment, Disposal, Reuse*. 3<sup>rd</sup> Edition. McGraw-Hill, Inc., New York.
7. Michigan Mercury Pollution Prevention Task Force. April 1996. *Mercury Pollution Prevention in Michigan: Summary of Current Efforts and Recommendations for Future Activities*.
8. Minnesota Pollution Control Agency, Michigan Department of Environmental Quality, and Wisconsin Department of Natural Resources. *The Waste Connection: Mercury in the Environment*.
9. MDEQ. 1994. *Wastewater Regulations for National Pollutant Discharge Elimination System (NPDES) Permits, Underground Injection Control (UIC) Permits, State Permits, Water Quality Based Effluent Limitations and Water Quality Certification*. Office of Pollution Control.
10. MDEQ. 2003. *State of Mississippi Water Quality Criteria for Intrastate, Interstate, and Coastal Waters*. Office of Pollution Control.
11. MDEQ. 2002. *Mississippi List of Waterbodies, Pursuant to Section 303(d) of the Clean Water Act*. Office of Pollution Control.
12. MDEQ. 1999. *Mississippi Water Quality Assessment, Federal Clean Water Act Section 305(b) Report*. Office of Pollution Control.

13. Tremblay, A., L. Cloutier, and M. Lucotte. 1998. Total mercury and methylmercury fluxes via emerging insects in recently flooded hydroelectric reservoirs and a natural lake. *Sci. Total Environ.* 219:209-221.
14. USEPA. 1997. Mercury study report to congress. EPA-452/R-97-003. Office of Air Quality, Planning and Standards. Office of Research and Development. Washington, DC.
15. USEPA. 1998. *Better Assessment Science Integrating Point and Nonpoint Sources, BASINS, Version 2.0 User's Manual*. U.S. Environmental Protection Agency, Office of Water, Washington, D.C.
16. USEPA. 1986. *Quality Criteria for Water 1986*. U.S. Environmental Protection Agency, Office of Water, Washington, D.C.
17. USEPA. Region 6. 2000. *Mercury TMDLs for Segments within Mermentau and Vermilion-Teche River Basins*. U.S. Environmental Protection Agency, Region 6, Dallas Texas.
18. Watras, C.J., K. A. Morrison, and R. C. Back. 1996. Mass balance studies of mercury and methylmercury in small, temperate/boreal lakes of the Northern Hemisphere. In” Baeyens, E., R. Ebinghaus, O. Vasiliev. (ed.) *Regional and global mercury cycles: sources, fluxes and mass balances*. Kluwer Academic Publ. Netherlands. Pp. 329-358.
19. Zillioux, E.J., D. B. Porcella, J. M. Benoit. 1993. Mercury cycling and effects in freshwater wetland ecosystems. *Environ. Toxicol. Chem.* 12:2245-2264.

## APPENDIX A: Fish Tissue Data

Fish Flesh Mercury Data in the Pascagoula River Watershed

| County             | Year | Site                                | Hg (ppm) |    | Species          | # Fish | Min. Wt. | Max. Wt. | Mean Wt. |
|--------------------|------|-------------------------------------|----------|----|------------------|--------|----------|----------|----------|
| Jackson            | 1996 | East Pascagoula River @ L.R. Marina | 1.31     | ** | Largemouth Bass  | 3      | 3.3      | 3.9      | 3.7      |
| Jackson            | 1996 | East Pascagoula River @ L.R. Marina | 0.88     |    | Largemouth Bass  | 4      | 1.3      | 2.1      | 1.6      |
| Jackson            | 1995 | East Pascagoula River @ L.R. Marina | 0.83     |    | Largemouth Bass  | 2      | 4.0      | 4.4      | 4.2      |
| Jackson            | 1995 | East Pascagoula River @ L.R. Marina | 0.9      |    | Largemouth Bass  | 5      | 0.9      | 1.5      | 1.1      |
| Jackson            | 1995 | East Pascagoula River @ L.R. Marina | ND       |    | Flathead Catfish | 2      | 4.2      | 5.5      | 4.9      |
| Jackson            | 1995 | East Pascagoula River @ L.R. Marina | ND       |    | Channel Catfish  | 1      | 6.8      | 6.8      | 6.8      |
|                    |      |                                     |          |    |                  |        |          |          |          |
| Jackson            | 1996 | West Pascagoula River Above I-10    | 1.84     | ** | Largemouth Bass  | 1      | 3.9      | 3.9      | 3.9      |
| Jackson            | 1996 | West Pascagoula River Above I-10    | 0.79     |    | Largemouth Bass  | 5      | 1.2      | 2.1      | 1.5      |
| Jackson            | 1996 | West Pascagoula River Above I-10    | 0.63     |    | Largemouth Bass  | 5      | 1.0      | 1.7      | 1.3      |
| Jackson            | 1996 | West Pascagoula River Above I-10    | 0.3      |    | Largemouth Bass  | 4      | 0.6      | 0.7      | 0.6      |
| Jackson            | 1997 | West Pascagoula River Above I-10    | 0.4      |    | Largemouth Bass  | 3      | 1.3      | 1.7      | 1.5      |
| Jackson            | 1997 | West Pascagoula River Above I-10    | 0.2      |    | Red Drum         | 2      | 8.0      | 10.0     | 9.0      |
| Jackson            | 1997 | West Pascagoula River Above I-10    | 0.61     |    | Largemouth Bass  | 2      | 2.3      | 2.8      | 2.6      |
| Jackson            | 1997 | West Pascagoula River Above I-10    | 0.52     |    | Largemouth Bass  | 5      | 1.0      | 1.3      | 1.1      |
| Jackson            | 1998 | West Pascagoula River Above I-10    | 0.6      |    | Largemouth Bass  | 4      | 1.5      | 2.0      | 1.7      |
| Jackson            | 1998 | West Pascagoula River Above I-10    | 0.4      |    | Sheepshead       | 2      | 2.9      | 3.5      | 3.2      |
| Jackson            | 1999 | West Pascagoula River Above I-10    | 0.1      |    | White Trout      | 2      | 0.4      | 0.5      | 0.5      |
| Jackson            | 1999 | West Pascagoula River Above I-10    | ND       |    | Atlantic Croaker | 5      | 0.2      | 0.3      | 0.2      |
| Jackson            | 2000 | West Pascagoula River Above I-10    | 0.11     |    | Gafftopsail      | 2      | 1.3      | 1.3      | 1.3      |
| Jackson            | 2000 | West Pascagoula River Above I-10    | ND       |    | Atlantic Croaker | 5      | 0.4      | 0.5      | 0.4      |
| Jackson            | 2001 | West Pascagoula River Above I-10    | 0.3      |    | Largemouth Bass  | 5      | 0.7      | 0.8      | 0.8      |
| Jackson            | 2001 | West Pascagoula River Above I-10    | 0.1      |    | Sheepshead       | 1      | 0.9      | 0.9      | 0.9      |
| ** Above 1.0 (ppm) |      |                                     |          |    |                  |        |          |          |          |



Fish Flesh Mercury Data in the Pascagoula River Watershed Cont'd

| County             | Year | Site                             | Hg (ppm) |    | Species          | # Fish | Min. Wt. | Max. Wt. | Mean Wt. |
|--------------------|------|----------------------------------|----------|----|------------------|--------|----------|----------|----------|
| Jackson            | 1996 | Pascagoula River @ Cumbest Bluff | 1.31     | ** | Largemouth Bass  | 2      | 2.7      | 4.6      | 3.7      |
| Jackson            | 1996 | Pascagoula River @ Cumbest Bluff | 0.77     |    | Bass sp.         | 5      | 1.1      | 1.3      | 1.3      |
| Jackson            | 1996 | Pascagoula River @ Cumbest Bluff | 1.06     | ** | Bowfin           | 2      | 5.1      | 5.9      | 5.5      |
|                    |      |                                  |          |    |                  |        |          |          |          |
| Jackson            | 1994 | Pascagoula River @ Poticaw Camp  | 1.15     | ** | Largemouth Bass  | 3      | 2.7      | 3.9      | 3.1      |
| Jackson            | 1994 | Pascagoula River @ Poticaw Camp  | 0.85     |    | Largemouth Bass  | 5      | 1.1      | 1.8      | 1.4      |
| Jackson            | 1995 | Pascagoula River @ Poticaw Camp  | 1.16     | ** | Largemouth Bass  | 1      | 3.6      | 3.6      | 3.6      |
| Jackson            | 1995 | Pascagoula River @ Poticaw Camp  | 0.87     |    | Largemouth Bass  | 5      | 1.5      | 2.1      | 1.8      |
| Jackson            | 1995 | Pascagoula River @ Poticaw Camp  | 0.58     |    | Largemouth Bass  | 4      | 0.6      | 1.0      | 0.8      |
| Jackson            | 1995 | Pascagoula River @ Poticaw Camp  | 1.44     | ** | Largemouth Bass  | 3      | 1.7      | 3.1      | 2.4      |
| Jackson            | 1995 | Pascagoula River @ Poticaw Camp  | 0.92     |    | Largemouth Bass  | 3      | 0.9      | 1.5      | 1.1      |
| Jackson            | 1995 | Pascagoula River @ Poticaw Camp  | 1.29     | ** | Flathead Catfish | 1      | 17.2     | 17.2     | 17.2     |
| Jackson            | 1995 | Pascagoula River @ Poticaw Camp  | 0.45     |    | Flathead Catfish | 3      | 3.7      | 5.0      | 4.3      |
| Jackson            | 1996 | Pascagoula River @ Poticaw Camp  | 1.73     | ** | Largemouth Bass  | 1      | 3.0      | 3.0      | 3.0      |
| Jackson            | 1996 | Pascagoula River @ Poticaw Camp  | 1.31     | ** | Largemouth Bass  | 1      | 2.4      | 2.4      | 2.4      |
| Jackson            | 1996 | Pascagoula River @ Poticaw Camp  | 1.1      | ** | Largemouth Bass  | 1      | 1.8      | 1.8      | 1.8      |
| Jackson            | 1996 | Pascagoula River @ Poticaw Camp  | 1.44     | ** | Largemouth Bass  | 1      | 1.3      | 1.3      | 1.3      |
| Jackson            | 1996 | Pascagoula River @ Poticaw Camp  | 0.98     |    | Largemouth Bass  | 1      | 1.3      | 1.3      | 1.3      |
| Jackson            | 1996 | Pascagoula River @ Poticaw Camp  | 0.68     |    | Largemouth Bass  | 1      | 0.9      | 0.9      | 0.9      |
| Jackson            | 1996 | Pascagoula River @ Poticaw Camp  | 0.97     |    | Largemouth Bass  | 1      | 0.9      | 0.9      | 0.9      |
| Jackson            | 1996 | Pascagoula River @ Poticaw Camp  | 0.59     |    | Largemouth Bass  | 1      | 0.9      | 0.9      | 0.9      |
| Jackson            | 1996 | Pascagoula River @ Poticaw Camp  | 0.58     |    | Largemouth Bass  | 1      | 0.8      | 0.8      | 0.8      |
| Jackson            | 1996 | Pascagoula River @ Poticaw Camp  | 0.98     |    | Largemouth Bass  | 1      | 0.7      | 0.7      | 0.7      |
| ** Above 1.0 (ppm) |      |                                  |          |    |                  |        |          |          |          |

Fish Flesh Mercury Data in the Pascagoula River Watershed Cont'd

| County             | Year | Site                        | Hg (ppm) |    | Species          | # Fish | Min. Wt. | Max. Wt. | Mean Wt. |
|--------------------|------|-----------------------------|----------|----|------------------|--------|----------|----------|----------|
| Jackson            | 1997 | Pascagoula River @ Road 614 | 0.95     |    | Bass sp.         | 2      | 2.1      | 2.7      | 2.4      |
| Jackson            | 1997 | Pascagoula River @ Road 614 | 0.53     |    | Flathead Catfish | 2      | 6.9      | 7.3      | 7.1      |
| Jackson            | 1997 | Pascagoula River @ Road 614 | 0.68     |    | Flathead Catfish | 1      | 12.9     | 12.9     | 12.9     |
| Jackson            | 1997 | Pascagoula River @ Road 614 | 1.13     | ** | Largemouth Bass  | 1      | 4.0      | 4.0      | 4.0      |
| Jackson            | 1997 | Pascagoula River @ Road 614 | 0.51     |    | Largemouth Bass  | 1      | 1.1      | 1.1      | 1.1      |
| Jackson            | 1997 | Pascagoula River @ Road 614 | 0.48     |    | Spotted Bass     | 1      | 1.3      | 1.3      | 1.3      |
|                    |      |                             |          |    |                  |        |          |          |          |
| George             | 1997 | Pascagoula River @ Benndale | 0.26     |    | Flathead Catfish | 3      | 3.6      | 4.5      | 4.2      |
| George             | 1997 | Pascagoula River @ Benndale | 0.17     |    | Blue Catfish     | 4      | 3.4      | 3.8      | 3.5      |
| George             | 1997 | Pascagoula River @ Benndale | 0.91     |    | Largemouth Bass  | 1      | 3.6      | 3.6      | 3.6      |
| George             | 1997 | Pascagoula River @ Benndale | 0.55     |    | Largemouth Bass  | 1      | 1.4      | 1.4      | 1.4      |
| George             | 1997 | Pascagoula River @ Benndale | 0.56     |    | Spotted Bass     | 1      | 0.7      | 0.7      | 0.7      |
| ** Above 1.0 (ppm) |      |                             |          |    |                  |        |          |          |          |

**APPENDIX B: Daily Load from NPDES Permitted Facilities**

| NPDES ID                | Facility Name  | Permitted Flow (MGD) | Mercury Criterion (µg/l) | Conversion Factor | Daily Load (g/day) |
|-------------------------|--|----------------------|--------------------------|-------------------|--------------------|
| MS0031828               | George County Schools<br>Central Elementary School               | 0.0160               | 0.012                    | 3.79              | 7.28E-04           |
| MS0044504               | Lucedale POTW  | 0.5000               | 0.012                    | 3.79              | 2.27E-02           |
| MS0020249               | MS Gulf Coast Regional Wastewater<br>Authority - Pascagoula POTW | 10.0000              | 0.012                    | 3.79              | 4.55E-01           |
| MS0028355               | Jackson County School District<br>East Central School            | 0.0500               | 0.012                    | 3.79              | 2.27E-03           |
| MS0028762               | Jackson County School District<br>Vancleave Junior High School   | 0.0240               | 0.012                    | 3.79              | 1.09E-03           |
| MS0032115               | Bluff Creek Mobile Home Park                                     | 0.0340               | 0.012                    | 3.79              | 1.55E-03           |
| MS0033375               | Kwik Kar Wash  | 0.0009               | 0.012                    | 3.79              | 4.09E-05           |
| MS0035637               | Gulf City Seafoods, Inc.   | 0.0297*              | 0.012                    | 3.79              | 1.35E-03           |
| MS0035769               | MDOT Interstate 10 Rest Area                                     | 0.0500               | 0.012                    | 3.79              | 2.27E-03           |
| MS0038326               | Jackson County School District<br>Vancleave High School          | 0.0165               | 0.012                    | 3.79              | 7.50E-04           |
| MS0042269               | Finicky Pet Food, Inc.   | 0.0040*              | 0.012                    | 3.79              | 1.82E-04           |
| MS0043010               | MS Gulf Coast Regional Wastewater<br>Authority - Gautier POTW    | 4.0000               | 0.012                    | 3.79              | 1.82E-01           |
| MS0043966               | Seachick, Inc.   | 4.4300*              | 0.012                    | 3.79              | 2.01E-01           |
| MS0046221               | Davis Laundry and Quickstop                                      | 0.0036               | 0.012                    | 3.79              | 1.64E-04           |
| MS0047007               | Circle V Traditions, Inc.  | 0.0024               | 0.012                    | 3.79              | 1.09E-04           |
| MS0048445               | Colle Towing Company, Inc.                                       | 0.0050               | 0.012                    | 3.79              | 2.27E-04           |
| MS0055379               | Pascagoula Water Treatment Plant -<br>Communny Avenue            | 60.4500*             | 0.012                    | 3.79              | 2.75E+00           |
| <b>ΣWLA<sub>s</sub></b> |  |                      |                          |                   | <b>3.62E+00</b>    |

\* This NPDES facility does not have a permit limit for flow. The facility is required to report flow information in each discharge monitoring report. The flow value given above is an average of the facility's historical flow data.

## APPENDIX C: Mercury Use Outline

### Sources of Mercury

#### I. Deliberate Use of Mercury

##### A. Use of Mercury for its Physical and Electrical Properties

1. Instruments
  - a. *Barometers*
  - b. *Hydrometers*
  - c. *Manometers*
  - d. *Pyrometers*
  - e. *Sphygmometers*
  - f. *Thermometers*
2. Lamps
  - a. *Fluorescent*
  - b. *High Pressure Sodium*
  - c. *Mercury Arc*
  - d. *Metal Halide*
  - e. *Neon*
  - f. *UV disinfectant*
3. Pivots
  - a. *WWTP Trickling Filter System*
  - b. *Lighthouses*
4. Switches
  - a. *Household Switches*
  - b. *Industrial Switches*
  - c. *Mercury Thermocouple*
  - d. *Tilt (Motion) Switches*
5. Electrical Equipment
  - a. *Rectifiers*
  - b. *Batteries {Including alkaline, button (Hg – Zn) and (Hg – Cd)}*
6. Toys and Games

##### B. Medical, Dental, and Veterinary Use

1. Pharmaceuticals
  - a. *Anesthetic*
  - b. *Antiseptic*
  - c. *Antineoplastic Agent*
  - d. *Antisyphilitic*
  - e. *Cathartic*
  - f. *Diuretic*
  - g. *Purgative*
2. Dental Amalgam
3. Disinfectant
  - a. *Phenyl Mercuric Acetate (PMA)*
  - b. *Thimerisol*
4. Diagnostic Reagents (see laboratory use)

C. Spiritist Use

1. Ingested, Dusted, Added to Bathing Solutions and Candles
  - a. *Asogue (Hg)*
  - b. *Precipitado Rojo (HgO)*
  - c. *Precipitado Amarillo (HgO)*
  - d. *Precipitado Blanco (Hg<sub>2</sub>Cl<sub>2</sub>)*

D. Laboratory Use

1. Slide Preparation
  - a. *Stain*
2. Electroanalysis
  - a. *Cathode*
3. Algae Sample Preservative
4. Reagents (used to analyze other chemicals)
  - a. *Acetic Acid*
  - b. *Acetone*
  - c. *Aldehyde*
  - d. *Ammonia*
  - e. *Arsenic*
  - f. *Barbital*
  - g. *Chloride*
  - h. *Chlorine*
  - i. *Citric Acid*
  - j. *CO in gas*
  - k. *Cystine*
  - l. *Glucose*
  - m. *HCN*
  - n. *Iron*
  - o. *Kjeldahl Nitrogen*
  - p. *Manganese*
  - q. *Mercury*
  - r. *Triophene*
  - s. *Vanadium*
  - t. *Wine Coloring*
  - u. *Zinc*

E. Mining/Metals Industry

1. Electrolysis
  - a. *Cathode*
2. Extracting Au and Ag from Ore
3. Extracting Au from Pb
4. Electroplating Al
5. Other Processes
  - a. *Etching Steel/Iron*
  - b. *Fire Gilding*
  - c. *Blackening Brass*

F. Chlor-Alkali Industry

1. Mercury Cell Process
  - a. *Production of Chlorine, Caustic Soda, Sodium Hydroxide and Products Manufactured with These Raw Materials*

G. Fungicide/Pesticide

1. Seed Protectant
2. Golf Courses
  - a. *Snow Mold Control*
3. Root Maggot Control
4. Imported Gray Goods (undyed textiles)
5. Paint and Glues
  - a. *Latex Paint\**
  - b. *Marine Paint\**
  - c. *Gold Porcelain Paint*
  - d. *Corrugated Cardboard Glue*

H. Preservative

1. Kyanizing Wood\*
2. Anatomical Specimens
3. Embalming\*
4. Tanning

I. Coloring

1. Pigment
  - a. *Colored Papers*
  - b. *Horn*
  - c. *Inks*
  - d. *Linen*
  - e. *Plastics*
  - f. *Rubber*
  - g. *Sealing Wax*
2. Stain for Wood\*
3. Mordant for Dye
  - a. *Beaver and Rabbit Pelts*

J. Other Deliberate Uses

1. Plastics
  - a. *Catalyst for Curing*
2. Fireworks
  - a. *Pharoah's Serpents and Bengal Green Lights*
3. Photography\*
  - a. *Intensifier*
  - b. *Magic Photograms*

**II. Production/Storage**

A. Mining

1. Mines with Mercury as the Primary Product
2. Mines with Secondary Production of Mercury

B. U.S. Federal Supply

C. Recycling

1. Facilities Include Fluorescent Lamp Recycling and Thermostat Recycling

### **III. By-Product/Contaminant**

#### **A. Combustion**

1. Incineration
  - a. *Municipal Solid Waste*
  - b. *Medical Waste*
  - c. *Sewage Sludge*
  - d. *Cremation*
2. Fuel Combustion
  - a. *Coal*
  - b. *Oil*
  - c. *Natural Gas*
  - d. *Wood*

#### **B. Vaporization**

1. Landfill Gas
2. Petroleum Refining
3. Wastewater Treatment Plants
4. Mining
  - a. *Smelting*
  - b. *Roasting*

#### **C. Product Contaminant**

1. Chloralkali Products

### **IV. Natural**

#### **A. Volcanoes**

#### **B. Mineralized Bedrock**

1. Cinnabar

Source: Michigan Mercury Pollution Prevention Task Force. April 1996. *Mercury Pollution Prevention in Michigan: Summary of Current Efforts and Recommendations for Future Activities.*

## APPENDIX D: Mississippi Fish Advisories

| MISSISSIPPI'S FISH TISSUE ADVISORIES<br>AND COMMERCIAL FISHING BANS<br>AUGUST 2001   |                   |             |   |
|--|-------------------|-------------|---|
| WATERBODY  | CHEMICAL          | DATE ISSUED | ACTION  |
| Little Conehoma Creek and Yockanookany River in Attala and Leake Counties. From Hwy 35 near Kosciusko, downstream to Hwy 429 near Thomastown   | PCB's             | June 1987   | Consumption Advisory<br>All Species<br>Commercial Fishing Ban   |
| Lake Susie, Oxbow Lake of Old Tallahatchie River in Panola County west of Batesville.  | PCB's             | Nov. 1989   | Same as above   |
| Escatawpa River from the Alabama state line to I-10.   | Mercury           | May 1995    | Limit Consumption Advisory for largemouth bass and large catfish (>27 in.)*   |
| Bogue Chitto River, entire length in Mississippi.  | Mercury           | May 1995    | Same as above   |
| Yockanookany River, entire length.   | Mercury           | May 1995    | Same as above   |
| Pearl River from Hwy 25 near Carthage, downstream to the Leake County Water Park.  | Mercury           | June 2001   | Same as above   |
| Enid Reservoir   | Mercury           | May 1995    | Same as above   |
| Yocona River from Enid Reservoir downstream to the confluence with the Tallahatchie River.   | Mercury           | Sept. 1996  | Same as above   |
| Pascagoula River, entire length.   | Mercury           | Sept. 1996  | Same as above   |
| Archusa Creek Water Park   | Mercury           | Sept. 1996  | Same as above   |
| Grenada Lake and Yalobusha River from the dam downstream to Holcomb.   | Mercury           | June 2001   | Same as above   |
| Pearl River from Hwy 25 near Carthage, downstream to the Leake County Water Park.  | Mercury           | June 2001   | Same as above   |
| Mississippi Delta - all waters from the mainline Mississippi River Levee on the West to the Bluff hills on the East.   | DDT,<br>Toxaphene | June 2001   | Limit Consumption Advisory for carp, buffalo, gar, and large catfish (>22 in.)****  |
| Roebuck Lake, LeFlore County   | DDT,<br>Toxaphene | June 2001   | Limit Consumption Advisory for carp, gar, and large catfish (>22 in.)**** No Consumption of Buffalo. Commercial Fishing Ban |
| Yazoo National Wildlife Refuge (all waters)  | DDT,<br>Toxaphene | 1975        | Closed to fishing**   |
| Gulf of Mexico   | Mercury           | May 1998    | King Mackerel<br><33" - no limit.<br>33-39" limit consumption.***<br>>39" - do not eat                                      |
| * The Mississippi State Health Department recommends that people limit the amount of bass and large catfish that they eat from these areas, because of high levels of mercury in the fish. Children under seven and women of child bearing age should eat no more than one meal of these fish every two months. Other adults should eat no more than one meal of these fish every two weeks. |                   |             |   |
| ** Precautionary advisory issued by U.S. Fish and Wildlife Service   |                   |             |   |
| *** The Mississippi State Health Department recommends that people limit the amount of 33-39" King Mackerel they eat from the Mississippi Gulf Coast. Children under seven and women of child bearing age should eat no more than one meal of these fish every two months. Other adults should eat no more than one meal of these fish every two weeks.                                      |                   |             |   |
| **** The Mississippi Department of Health recommends that people limit their consumption of these fish to no more than one meal every two weeks.   |                   |             |   |